

University of Mississippi

eGrove

Faculty and Student Publications

Chemistry and Biochemistry

1-1-2019

Teleneurology service provided via tablet technology: 3-year outcomes and physician satisfaction

Kelly Harper

Vanderbilt University Medical Center

Megan McLeod

University of Mississippi

Summer K. Brown

Vanderbilt University

Georgia Wilson

Vanderbilt University Medical Center

Maxim Turchan

Vanderbilt University Medical Center

See next page for additional authors

Follow this and additional works at: https://egrove.olemiss.edu/chem_facpubs

Recommended Citation

Harper, K., McLeod, M., Brown, S., Wilson, G., Turchan, M., Gittings, E., Riebau, D., Baker, M., Zimmerman, E., & Charles, D. (2019). Teleneurology service provided via tablet technology: 3-year outcomes and physician satisfaction. *Rural and Remote Health*. <https://doi.org/10.22605/RRH4743>

This Article is brought to you for free and open access by the Chemistry and Biochemistry at eGrove. It has been accepted for inclusion in Faculty and Student Publications by an authorized administrator of eGrove. For more information, please contact egrove@olemiss.edu.

Authors

Kelly Harper, Megan McLeod, Summer K. Brown, Georgia Wilson, Maxim Turchan, Emily M. Gittings, Derek Riebau, Mark Baker, Eli Zimmerman, and David Charles



PROJECT REPORT

Teleneurology service provided via tablet technology: 3-year outcomes and physician satisfaction

AUTHORS

Kelly Harper¹ BPS, Program Manager



Megan McLeod² BBA, Medical Student



Summer K Brown³ BA, Masters student



Georgia Wilson⁴ BA, Clinical Research Coordinator I *

Maxim Turchan⁵ BA, Health and Policy Services Analyst II

Emily M Gittings⁶ Student

Derek Riebau⁷ MD, Assistant Professor



Mark Baker⁸ MD, Assistant Professor



Eli Zimmerman⁹ MD, Assistant Professor



David Charles¹⁰ MD, Professor and Vice-Chair of Neurology, Medical Director Vanderbilt TeleHealth

CORRESPONDENCE

*Ms Georgia Wilson georgia.j.wilson@vanderbilt.edu

AFFILIATIONS

^{1, 5, 7, 8, 9, 10} Vanderbilt University Medical Center, Nashville, Tennessee, USA

² University of Mississippi, Oxford, Mississippi, USA

³ Vanderbilt University, Nashville, Tennessee, USA

⁴ Department of Neurology, Vanderbilt University Medical Center, Nashville, Tennessee, USA

⁶ McGill University, Montreal, Canada

PUBLISHED

4 March 2019 Volume 19 Issue 1

HISTORY

RECEIVED: 25 January 2018

REVISED: 31 October 2018

ACCEPTED: 20 November 2018

CITATION

Harper K, McLeod M, Brown SK, Wilson G, Turchan M, Gittings EM, Riebau D, Baker M, Zimmerman E, Charles D. Teleneurology service provided via tablet technology: 3-year outcomes and physician satisfaction. *Rural and Remote Health* 2019; 19: 4743. <https://doi.org/10.22605/RRH4743>

Except where otherwise noted, this work is licensed under a [Creative Commons Attribution 4.0 International Licence](https://creativecommons.org/licenses/by/4.0/)

ABSTRACT:

Introduction: This study aimed to demonstrate that teleneurology consultations conducted via tablet technology are an efficient and cost-effective means of managing acute neurologic emergencies at community-based hospitals and that utilizing such technology yields high community physician satisfaction.

Method: During a 39-month period, Vanderbilt University Medical Center in Tennessee USA, provided teleneurology services to 10 community-based hospitals that lacked adequate neurology coverage. Hospitalists at one community-based hospital were not comfortable treating any patient with a neurologic symptom, resulting in 100% of those patients being transferred. This facility now retains more than 60% of neurology patients. For less than US\$1200, these hospitals were able to meet the only capital expenditure required to launch this service: the purchase of handheld tablet computers. Real-time teleneurology consultations were conducted via tablet using two-way video conferencing, radiologic image sharing, and medical record documentation.

Keywords:

neurology, seizure, stroke, telehealth, telemedicine, USA.

Community physicians were regularly surveyed to assess satisfaction.

Results: From February 2014 to May 2017, 3626 teleneurology consultations were conducted. Community physicians, in partnership with neurologists, successfully managed 87% of patients at the community-based hospital. Only 13% of patients required transfer to another facility for a higher level of care. The most common diagnoses included stroke (34%), seizure (11%), and headache/migraine (6%). The average time for the neurologist to answer a request for consultation page and connect with the community physician was 10.6 minutes. Ninety-one percent of community physicians were satisfied or somewhat satisfied with the overall service.

Conclusion: In the assessment of neurology patients, tablets are a more cost-effective alternative to traditional telehealth technologies. The devices promote efficiency in consultations through ease of use and low transfer rates, and survey results indicate community physician satisfaction.

FULL ARTICLE:

Introduction

According to the US Department of Health and Human Services, 67.8 million Americans live in a primary medical health professional shortage area¹. Telemedicine – the provision of medical care remotely using telecommunications technology – is increasingly being utilized to address this shortage of healthcare professionals. Dating from the late 1960s in the USA, telemedicine began with the ‘initial use of the closed-circuit, television system’². The practice has evolved to include technologies like telemedicine carts and more modern devices, such as tablets and smartphones. Ultimately, these telemedicine advancements enable physicians and other healthcare providers to efficiently and effectively increase access to specialty care for individuals that are ‘medically underserved, resulting in improved clinical outcomes, reduced costs’, and fewer unnecessary patient transfers³⁻⁸.

Disparities in access to health care are immense for specialties such as neurology. For large rural populations, the ratio of neurologists to patients is approximately 1:30 000⁹. The abundant need for neurological care is evident in the millions of individual visits to emergency departments each year for conditions like seizures and migraine-related headaches^{10,11}. In addition, stroke is classified as the fifth leading cause of death in the USA, and treatment by a neurologist following a stroke is demonstrated to improve clinical outcomes^{12,13}. The demand for neurological care far outweighs its accessibility; thus, the dawn of telemedicine has given rise to teleneurology.

Teleneurology employs telehealth technologies to serve patients with a variety of neurological conditions in both emergent and non-emergent settings^{14,15}. It is associated with effectiveness,

patient satisfaction, and a reduction in both time and money^{16,17}. In teleneurology assessments of 354 veterans with a variety of neurological conditions, Davis et al. (2017) found ‘average time savings of 5 h and 325 miles [523 km] driven, plus at least [US]\$48,000 total cost savings’ when compared to in-person assessments¹⁶.

Despite its cost-saving and time-saving benefits, the practice of teleneurology in hospitals is often discouraged by the price of its associated technologies and a lack of familiarity with the equipment¹⁸⁻²⁰. The most prevalent device in telemedicine is a large rolling cart equipped with a computer, camera, and screen¹⁹. Traditional telemedicine carts typically cost between US\$5000 and US\$25,000, sometimes incurring further additional costs due to annual service contracts. Not only are these outdated technologies costly, but physicians often poorly understand how to utilize them²⁰. High costs, difficulty of use, and the training required for standard telemedicine technology hinders its cost-effectiveness and efficiency.

To test the effectiveness of a less expensive technology, Anderson et al. (2013) performed teleneurology assessments of stroke patients using an iPhone 4. For most items on the National Institutes of Health Stroke Scale, researchers found near-perfect agreement between in-person and iPhone 4 consultations, and they concluded that the device is a time-saving, cost-effective alternative to typical telemedicine technologies¹⁸. A similar study employing iPads and the two-way live audio/video application FaceTime to conduct pediatric assessments demonstrated that the handheld technology was not only thousands of dollars cheaper than standard telemedicine equipment, but all healthcare

providers found the devices easy to use²¹.

To evaluate the benefits of similar technologies, the Department of Neurology at Vanderbilt University Medical Center (VUMC) in Tennessee, USA, introduced a teleneurology consultation service in February 2014. Using three software applications on low-cost, handheld tablet technology, teleneurologists are able to examine patients in community-based hospitals, view medical images, and record their clinical documentation, including recommendations, in patients' electronic medical records³.

Method

A total of 3626 teleneurology consultations were conducted over the 39-month period of 1 February 2014 – 31 May 2017. Patients ranged in age from 18 to 107 years (63+18 years, 55% female). Additionally, at the beginning of each month, physicians from the 10 community-based hospitals who called a teleneurology consult during the previous month were emailed a link and invited to complete a satisfaction survey based on their experiences during that time frame.

To conduct consultations, teleneurologists and community-based physicians were equipped with iPads and FaceTime. The teleneurologists' tablets also contained Jenesis and StarPanel, closed-group applications for radiologic image sharing and medical record documentation, respectively. The total cost of the tablet and software was US\$1200 for each community hospital. Information technology and risk assessment staff at VUMC determined that the security of these systems complied with *Health Insurance Portability and Accountability Act* (HIPAA) standards when linked over encrypted and password-protected internet connection³. Communications were protected using authentication and encryption provided by FaceTime. Password-protected, WPA2 Enterprise wireless sessions were used to authenticate and encrypt local traffic³.

Jenesis is used by community-based hospitals to upload and share radiologic images with teleneurologists. This application is hosted at VUMC, receives images from client imaging systems, and provides temporary storage of images until they are reviewed by the teleneurologist³. The StarPanel application is an integrated interface to the VUMC clinical information system that enabled secure access to the patient's medical record where teleneurology consultation notes were composed, stored, and transmitted to community-based hospitals for inclusion in their electronic medical records system³.

VUMC staff recorded the time of the initial consult request, the time at which the on-call neurologist was paged, and the time at which the neurologist and physician at the community-based hospital were connected. Study data were collected and managed using REDCap electronic data capture tools. REDCap (Research Electronic Data Capture) is a secure, web-based application

designed to support data capture for research studies, providing (1) an intuitive interface for validated data entry, (2) audit trails for tracking data manipulation and export procedures, (3) automated export procedures for seamless data downloads to common statistical packages, and (4) procedures for importing data from external sources²².

Results

A total of 3626 teleneurology consultations were conducted. In partnership with teleneurologists, the community physicians were able to successfully manage 87% of patients at their facility. Only 13% required transfer to another facility for a higher level of care. The most common diagnoses included stroke (34%), seizure (11%), and headache/migraine (6%).

Table 1 details the time required to establish a teleneurology connection. The average time between the community-based physician's initial request and the access center paging the on-call teleneurologist was 10.4 minutes and the average time between paging the teleneurologist and establishing a connection with the community physician was 10.6 minutes. For both of these times, the median was 8 minutes. The average time between the initial call for a consultation and connection to the teleneurologist and community physician was 21 minutes and the median was 17 minutes. Most consultations (75%) occurred between 9 am and 8 pm (Fig1).

Physician satisfaction surveys asked community physicians to rate the timeliness of connecting with the teleneurologist, the timeliness of receiving clinical documentation, whether they found the teleneurology service to be helpful in managing the care of patients, and their overall satisfaction with the service during a given month. After noting an insufficient initial response rate, this survey was redesigned, applying advanced survey methodologies including language brevity, simplified wording, and relevance to the participant²³⁻²⁶. In addition, survey invitations and reminders were sent at optimal times to generate a response^{27,28}. Data from the physician satisfaction surveys indicated that community physicians had an overwhelmingly positive experience with the teleneurology service (Table 2). Ninety-one percent reported being satisfied or somewhat satisfied with the overall service, 3% felt neutrally, and 6% were dissatisfied. It is worth noting that instances of dissatisfaction were not due to insufficient care nor did they result in harm to any patient. Most community physicians (95%) were satisfied or somewhat satisfied with the timeliness of connecting with the teleneurologist and the majority (96%) were also satisfied or somewhat satisfied with the helpfulness of the teleneurologist in managing patient care. Fewer community physicians (85%) were satisfied with the speed of documentation arrival after completion of a consultation. The response rate to these satisfaction surveys was initially low at 11% (69/628), but was increased to 17% (38/224) following the survey redesign, detailed above, that was implemented in December 2016.

Table 1: Mean and median times to establish teleconnection with neurologists during community physician consultations (N=3626)

Request/notification	Mean time (min)	Median time (min)
Initial request to neurologist notification	10.4	8
Neurologist notification to connection with community physician	10.6	8
Initial request to connection with community physician	21	17

Table 2: Community physician satisfaction with teleneurology (N=3626)

Factor	Satisfaction rating		
	Very satisfied/somewhat satisfied (%)	Neutral (%)	Somewhat dissatisfied/very dissatisfied (%)
Timeliness of return call	95	0	5
Patient care management	96	0	4
Speed of documentation	85	9	6
Service overall	91	3	6




Figure 1: Teleneurology consult volume, by hour.

Discussion

VUMC’s teleneurology service not only appeals to community physicians, but is also an efficient and cost-effective means of providing urgent neurologic consultation to community-based hospitals. For these hospitals, the cost of the tablet technology (less than US\$1200) is considerably less than the cost of outdated traditional telemedicine carts (US\$4999). The use of tablets also promotes efficiency in their ease of use and community healthcare providers’ existing familiarity with such devices.

These results indicate that a large majority of patients at community-based hospitals needing emergent neurologic care are able to be successfully treated via tablet technology using HIPAA-compliant two-way video conferencing, radiologic image-sharing, and electronic medical record capabilities. Tablet technology is accessible, portable, and has low implementation and operational costs. A high proportion of community physicians expressed satisfaction with the teleneurology service provided via this technology platform. Although fewer were satisfied with the timeliness of documentation return, this is likely a result of differing workflows between hospitals for the entry of forwarded medical record documentation.

Tertiary academic healthcare centers often operate at near-capacity while also serving as level one trauma centers²¹. Therefore, it is essential to leverage technology to project specialty care to community-based hospitals so that patients may be treated close to home and only transferred when they require

medical services or procedures uniquely available at academic healthcare centers.

The high level of community physician satisfaction and low rate of patient transfer noted in this analysis are consistent with other studies on the use of telemedicine to provide specialty care to rural communities; Marcin and associates reported a 97.5% community provider satisfaction rate with the overall quality of pediatric subspecialty telemedicine consultations²⁹. Latifi and associates found that telemedicine consultations significantly reduced the need for transfer of trauma patients and improved quality of treatment³⁰.

This study used the metrics of patient transfer, time to establish connection, and community physician satisfaction to evaluate a teleneurology program. Additional studies are needed to analyze patient satisfaction and clinical outcomes of services provided using teleneurology. Future research should also aim to address the limitations of this study. This could include comparing wait times of standard in-person neurology consultations and teleneurology consultations.

The use of telehealth in the provision of specialty and subspecialty care to rural communities is promising. Based on the early results from this program, VUMC intends to expand its telehealth services to other community-based hospitals throughout south-eastern USA and the Vanderbilt Health Affiliated Network. Effective telehealth services are an essential element of successful population health programs and they allow community-based

physicians and hospitals to effectively manage the care of patients in need of emergent specialty care, reducing burdens on the healthcare system, patients, and their support systems.

Author disclosure: Vanderbilt University Medical Center receives

income from Allergan, Ipsen, Medtronic, Merz, USWorldmeds, Boston Scientific, and St Jude for research or educational programs led by Dr Charles. Dr Charles receives income from Allergan, Ipsen and Revance for consulting services.

REFERENCES:

- 1** US Department of Health and Human Services, Bureau of Health Workforce Health Resources and Services Administration (HRSA). Designated health professional shortage area statistics. Available: <https://datawarehouse.hrsa.gov/Tools/HDWReports/Reports.aspx> (Accessed 28 July 2016).
- 2** Darkins AW, Cary MA. *Telemedicine and telehealth: principles, policies, performance, and pitfalls*. Springer Publishing Company, 2000; 6.
- 3** Cunningham PA, Harper K, Hermann L, Riebau, D, Humphrey A, Meyers H, et al. Teleneurology consultations for acute neurologic emergencies. *Tennessee Medicine E-Journal* 2015; **1(4)**.
- 4** Bynum AB, Irwin CA. Evaluation of the effect of consultant characteristics on telemedicine diagnosis and treatment. *International Journal of Telemedicine and Applications* 2011; **3**: 1-7.
- 5** Schwamm LH, Rosenthal ES, Hirshberg A, Schaefer PW, Little EA, Kvedar JC, et al. Virtual TeleStroke support for the emergency department evaluation of acute stroke. *Academic Emergency Medicine* 2004; **11**: 1193-1197. <https://doi.org/10.1197/j.aem.2004.08.014> PMID:15528584
- 6** Marcin JP, Nesbitt TS, Cole SL, Knuttel RM, Hilty DM, Prescott PT, et al. Changes in diagnosis, treatment, and clinical improvement among patients receiving telemedicine consultations. *Telemedicine Journal and e-Health* 2005; **11(1)**: 36-43. <https://doi.org/10.1089/tmj.2005.11.36> PMID:15785219
- 7** Aoki N, Dunn K, Johnson-Throop KA, Turley JP. Outcomes and methods in telemedicine evaluation. *Telemedicine and e-Health* 2003; **9(4)**: 393-401. <https://doi.org/10.1089/153056203772744734> PMID:14980098
- 8** Gillum LA, Johnston SC. Influence of physician specialty on outcomes after acute ischemic stroke. *Journal of Hospital Medicine* 2008; **3**: 184-192. <https://doi.org/10.1002/jhm.313> PMID:18570345
- 9** Association of American Medical Colleges. State physician workforce data book. 2015. Available: <https://www.aamc.org/data/workforce/reports/442830/statedataandreports.html> (Accessed 15 October 2017).
- 10** Pallin DJ, Goldstein JN, Moussally JS, Pelletier AJ, Green AR, Camargo CA. Seizure visits in US emergency departments: epidemiology and potential disparities in care. *International Journal of Emergency Medicine* 2008; **1(2)**: 97-105. <https://doi.org/10.1007/s12245-008-0024-4> PMID:19384659
- 11** Lucado J, Paez K, Elixhauser A. *Headaches in U.S. hospitals and emergency departments, 2008*. HCUP statistical brief #111. 2011. Available: <http://www.hcup-us.ahrq.gov/reports/statbriefs/sb111.pdf> (Accessed 25 July 2017).
- 12** Xu JQ, Murphy SL, Kochanek KD, Arias E. *Mortality in the United States, 2015*. 2016 NHCS data brief no. 267. Available: <https://www.cdc.gov/nchs/products/databriefs/db267.htm> (Accessed 15 October 2017).
- 13** Goldstein L, Matchar DB, Hoff-Lindquist J, Samsa GP, Horner RD. VA Stroke Study: neurologist care is associated with increased testing but improved outcomes. *Neurology* 2003; **61**: 792-796. <https://doi.org/10.1212/01.WNL.0000082724.77447.3A> PMID:14504322
- 14** Patterson V. Teleneurology. *Journal of Telemedicine and Telecare* 2005; **11(2)**: 55-59. <https://doi.org/10.1258/1357633053499840> PMID:15829048
- 15** Wechsler LR, Tsao JW, Levine SR, Swain-Eng RJ, Adams RJ, Demaerschalk BM, et al. Teleneurology applications: report of the telemedicine work group of the American Academy of Neurology. *Neurology* 2013; **80**: 670-676. <https://doi.org/10.1212/WNL.0b013e3182823361> PMID:23400317
- 16** Davis LE, Coleman J, Harnar J, King MK. Teleneurology: successful delivery of chronic neurologic care to 354 patients living remotely in a rural state. *Telemedicine Journal and e-Health* 2014; **20**: 473-477. <https://doi.org/10.1089/tmj.2013.0217> PMID:24617919
- 17** Beck CA, Beran DB, Biglan KM, Boyd CM, Dorsey ER, Schmidt PN, et al. National randomized controlled trial of virtual house calls for Parkinson disease. *Neurology* 2017; **89(11)**: 1152-1161. <https://doi.org/10.1212/WNL.0000000000004357> PMID:28814455
- 18** Anderson ER, Smith B, Ido M, Frankel M. Remote assessment of stroke using the iPhone 4. *Journal of Stroke and Cerebrovascular Diseases* 2013; **22(4)**: 340-344. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2011.09.013> PMID:22018507
- 19** Weschler LR. Advantages and limitations of teleneurology. *JAMA Neurology* 2015; **72(3)**: 349-354. <https://doi.org/10.1001/jamaneurol.2014.3844> PMID:25580942
- 20** Ayatollahi H, Sarabi FZP, Langaizadeh M. Clinicians' knowledge and perception of telemedicine technology. *Perspectives in Health Information Management* 2015; **12**.
- 21** Patel S, Hertzog JH, Penfil S, Slamon N. A prospective pilot study of the use of telemedicine during pediatric transport: a high-quality, low-cost alternative to conventional telemedicine systems. *Pediatric Emergency Care* 2015; **31(9)**: 611-616. <https://doi.org/10.1097/PEC.0000000000000544> PMID:26335229
- 22** Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap) – a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics* 2009; **42(2)**: 377-381. <https://doi.org/10.1016/j.jbi.2008.08.010>

PMid:18929686

23 Flanigan T, McFarlane ES, Cook S. Conducting survey research among physicians and other medical professionals – a review of current literature. In: *Proceedings of American Association for Public Opinion Research*. New Orleans, LA: American Association for Public Opinion Research, 2008.

24 Thomas M, Rogers R, Maclean R. Collecting data from physicians via web-based surveys: recommendations for improving response rates. *The Internet Journal of Medical Informatics* 2002; **1(1)**.

25 Office of Quality Improvement, University of Wisconsin-Madison. Survey fundamentals. 2010. Available: https://oqi.wisc.edu/resourcelibrary/uploads/resources/Survey_Guide.pdf (Accessed 24 October 2016).

26 Thorpe C, Ryan B, McLean SL, Burt A, Stewart M, Brown JB, et al. How to obtain excellent response rates when surveying physicians. *Family Practice* 2009; **26**: 65-68. <https://doi.org/10.1093/fampra>

/cmn097 PMid:19074758

27 Van Dessel G. What's the best time to send a survey? 2015. Available: <https://www.checkmarket.com/blog/survey-invitations-best-time-send/> (Accessed 1 November 2016).

28 Zheng J. What day of the week should you send your survey? 2011. Available: <https://www.surveymonkey.com/curiosity/day-of-the-week/> (Accessed 24 October 2016).

29 Marcin JP, Ellis J, Mawis R, Nagrampa E, Nesbitt TS, Dimand RJ. Using telemedicine to provide pediatric subspecialty care to children with special health care needs in an underserved rural community. *Pediatrics* 2004; **113(1)**: 1-6. <https://doi.org/10.1542/peds.113.1.1> PMid:14702439

30 Latifi R, Hadeed GJ, Rhee P, O'Keeffe T, Friese RS, Wynne JL, et al. Initial experiences and outcomes of telepresence in the management of trauma and emergency surgical patients. *American Journal of Surgery* 2009; **6(1)**: 905-910. <https://doi.org/10.1016/j.amjsurg.2009.08.011> PMid:19969150

This PDF has been produced for your convenience. Always refer to the live site <https://www.rrh.org.au/journal/article/4743> for the Version of Record.