

University of Mississippi

eGrove

Faculty and Student Publications

Pharmacy, School of

5-1-2019

A New Prenylated Isoflavonoid from *Limonium leptophyllum*

Asset Dizamatova

Al Farabi Kazakh National University

Kamila Zhumanova

Al Farabi Kazakh National University

Galiya E. Zhusupova

Al Farabi Kazakh National University

Aizhan I. Zhussupova

Al Farabi Kazakh National University

Radhakrishnan Srivedavyasari

University of Mississippi School of Pharmacy

See next page for additional authors

Follow this and additional works at: https://egrove.olemiss.edu/pharmacy_facpubs



Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)

Recommended Citation

Dizamatova, A., Zhumanova, K., Zhusupova, G. E., Zhussupova, A. I., Srivedavyasari, R., Ibrahim, M. A., & Ross, S. A. (2019). A New Prenylated Isoflavonoid From *Limonium leptophyllum*. *Natural Product Communications*, 14(5), 1934578X1984413. <https://doi.org/10.1177/1934578X19844137>

This Article is brought to you for free and open access by the Pharmacy, School of at eGrove. It has been accepted for inclusion in Faculty and Student Publications by an authorized administrator of eGrove. For more information, please contact egrove@olemiss.edu.

Authors

Asset Dizamatova, Kamila Zhumanova, Galiya E. Zhusupova, Aizhan I. Zhussupova, Radhakrishnan Srivedavyasari, Mohamed A. Ibrahim, and Samir A. Ross

A New Prenylated Isoflavonoid From *Limonium leptophyllum*

Natural Product Communications
 May 2019: 1–5
 © The Author(s) 2019
 Article reuse guidelines:
sagepub.com/journals-permissions
 DOI: 10.1177/1934578X19844137
journals.sagepub.com/home/npx



Asset Dizamatova¹, Kamila Zhumanova¹,
 Galiya E. Zhusupova¹, Aizhan I. Zhussupova¹, Radhakrishnan Srivedavyasari²,
 Mohamed A. Ibrahim^{2,3}, and Samir A. Ross^{2,4}

Abstract

Phytochemical investigation of methanolic extract of *Limonium leptophyllum* (Plumbaginaceae), led to the isolation of 1 new isoflavonoid with a rare 5-membered dihydrofuran ring (**1**, leptoisoflavone A) and 8 known compounds. The known isolated compounds were identified as euchrenone b₉ (**2**), auricularin (**3**), kaempferol (**4**), avicularoside (**5**), myricetin-3-arabinoside (**6**), *trans*-*N*-feruloyltyramine (**7**), *trans*-*N*-caffeoyltyramine (**8**), and β-sitosterol (**9**). The crude methanolic extract exhibited moderate activity toward endocannabinoid receptors. Auricularin (**3**) showed activity toward cannabinoid receptor type 1 (86.7% displacement with IC₅₀ 8.92 μM).

Keywords

Limonium leptophyllum, isoflavonoids, leptoisoflavone A, isolation, cannabinoid receptors

Received: October 12th, 2018; Accepted: January 7th, 2019.

The genus *Limonium* (Plumbaginaceae family) is known as halophytes and consists of 300 species worldwide, of which 18 are found in Kazakhstan.¹ Species in this genus are also known as statice, caspia, sea-lavender, or marsh-rosemary. The genus is widely distributed in Asia, Africa, Australia, Europe, and North America. More recently, several species of the *Limonium* genus are used to treat arthritis and fever.² This genus possesses anti-inflammatory, antibacterial,^{3,4} antiviral,⁵ and cytotoxic properties.³ Previous phytochemical and biological studies on the *Limonium* genus have shown the presence of different classes of compounds with various biological activities.^{6–14} In continuation of our previous work and search for novel plant-derived biological agents,^{6,7,15–17} further investigations on the phytochemical constituents of *Limonium leptophyllum* (Schrenk) O. Kuntze were carried out. Herein, we are reporting the isolation and characterization of 1 new and 8 known compounds from *L. leptophyllum*.

Compound **1** was obtained as a yellow solid. The high-resolution mass spectrometric (MS) data showed a peak corresponding to the molecular formula C₂₅H₂₂O₅, based on the [M + H]⁺ ion signal at *m/z* 403.1532 (calc. 403.1545). The ¹³C nuclear magnetic resonance (NMR) data of **1** (Table 1) showed resonances of 25 carbon atoms, which were classified by distortionless enhancement by polarization transfer 135 and heteronuclear single quantum coherence (HSQC) experiments as 3 methyls, 2

methylenes, 8 methines, and 12 quaternary carbons. The ¹H and ¹³C NMR spectra (Table 1) showed resonances of 3 methyls [$\delta_{\text{H}}/\delta_{\text{C}}$ 1.48/28.3, 1.52/28.5, and 1.92/19.9 (C-5''', 4''', and 5'')], methylene [$\delta_{\text{H}}/\delta_{\text{C}}$ 3.16 (dd, *J* = 9.0, 14.0 Hz), 3.03 (dd, *J* = 4.5, 14.0 Hz)/23.4 (C-1'')], an oxymethine proton [$\delta_{\text{H}}/\delta_{\text{C}}$ 4.45 (dd, *J* = 4.5, 9.0 Hz)/86.6 (C-2'')], olefinic methylene [$\delta_{\text{H}}/\delta_{\text{C}}$ 5.01 (s)/113.1 (C-4'')], a pair of olefinic protons [$\delta_{\text{H}}/\delta_{\text{C}}$ 5.61 (d, *J* = 10.0 Hz)/127.3, 6.71 (d, *J* = 10.0 Hz)/115.0, (C-2''', 1''')], a set of A₂B₂/ortho coupled protons of 1,4-disubstituted aromatic ring [$\delta_{\text{H}}/\delta_{\text{C}}$ 6.90 (d, *J* = 8.5 Hz)/115.7 (C-3', 5'), and 7.39 (d, *J* = 8.5 Hz)/130.5 (C-2', 6')], and an oxygenated aromatic proton [$\delta_{\text{H}}/\delta_{\text{C}}$ 7.90 (s)/152.7 (C-2)]. Furthermore, the ¹³C NMR showed

¹ Faculty of Chemistry and Chemical Technology, Al-Farabi Kazakh National University, Almaty, Republic of Kazakhstan

² National Center for Natural Products Research, University of Mississippi, USA

³ Department of Chemistry of Natural Compounds, National Research Center, Cairo, Egypt

⁴ Department of BioMolecular Sciences, Division of Pharmacognosy, School of Pharmacy, University of Mississippi, USA

Corresponding Author:

Samir A. Ross, Department of BioMolecular Sciences, Division of Pharmacognosy, School of Pharmacy, University of Mississippi, University, MS 38677, USA.

Email: ross@olemiss.edu



Table 1. ^1H and ^{13}C NMR Data for Compounds **1** and **2** in CDCl_3 (δ_{C} and δ_{H} in parts per minute; J in Hertz).

Position	Leptoisoflavone A (1)		Euchrenone b ₉ (2)	
	^1H NMR ^a	^{13}C NMR ^b	^1H NMR ^a	^{13}C NMR ^b
2	7.90 (s)	152.7 (D)	7.87 (s)	152.58 (D)
3	-	127.3 (S)	-	123.05 (S)
4	-	181.2 (S)	-	181.10 (S)
4a	-	105.6 (S)	-	105.7 (S)
5	-	159.9 (S)	-	160.09 (S)
6	-	109.6 (S)	-	110.03 (S)
7	-	157.5 (S)	-	157.44 (S)
8	-	101.1 (S)	-	100.99 (S)
8a	-	151.1 (S)	-	150.8 (S)
1'	-	123.7 (S)	-	123.63 (S)
2'	7.39 (d, 8.5)	130.5 (D)	7.37 (d, 8.0)	130.39 (D)
3'	6.90 (d, 8.5)	115.7 (D)	6.85 (d, 8.0)	115.70 (D)
4'	-	156.1 (S)	-	156.16 (S)
5'	6.90 (d, 8.5)	115.7 (D)	6.85 (d, 8.0)	115.70 (D)
6'	7.39 (d, 8.5)	130.5 (D)	7.37 (d, 8.0)	130.39 (D)
1''	3.16 (dd, 9.0, 14.0)	23.4 (T)	2.90 (dd, 9.2, 14.0)	29.20 (T)
	3.03 (dd, 4.5, 14.0)		3.03 (dd, 3.6, 14.0)	
2''	4.45 (dd, 4.5, 9.0)	86.6 (D)	4.30 (dd, 3.6, 9.2)	76.14 (D)
3''	-	144.4 (S)	-	147.57 (S)
4''	5.01 (s)	113.1 (T)	4.85 (s), 5.03 (s)	110.40 (T)
5''	1.92 (s)	19.9 (Q)	1.88 (s)	18.39 (Q)
1'''	6.71 (d, 10.0)	115.0 (D)	5.60 (d, 10.0)	127.22 (D)
2'''	5.61 (d, 10.0)	127.3 (D)	6.71 (d, 10.0)	114.99 (D)
3'''	-	78.8 (S)	-	78.60 (S)
4'''	1.52 (s)	28.5 (Q)	1.51 (s)	28.57 (Q)
5'''	1.48 (s)	28.3 (Q)	1.45 (s)	28.37 (Q)

NMR, nuclearmagnetic resonance.

Recorded at ^a500 MHz, ^b125MHz.

characteristic resonances attributed to an isoflavonoid skeleton of the conjugated oxo group [δ_{C} 181.2 (C-4)], 5 nonprotonated aromatic carbons [δ_{C} 101.1, 105.6, 109.6, 123.7, and 127.3 (C-8, 4a, 6, 1', and 3)], 4 oxygenated aromatic carbons [δ_{C} 159.9, 157.5, 156.1, and 151.1 (C-5, 7, 4', and 8a)], nonprotonated olefinic carbon [δ_{C} 144.4 (C-3'')], and nonprotonated oxygenated carbon [δ_{C} 78.8 (C-3''')]. The assignments of ^1H and ^{13}C NMR spectroscopic data of **1** (Table 1) were based on HSQC, heteronuclear multiple bond correlation (HMBC) correlations, and ^1H - ^1H correlated spectroscopy (COSY) couplings (Figure 1). The HMBC spectrum of compound **1** showed the following key 2J and 3J correlations (Figure 1): H-4''' and H-5''' exhibited correlation with C-2''', C-3''', and with each other, which confirmed the attachment of methyls to C-3'''. The methyl (H-5'') in the isoprenyl moiety showed correlations with C-2'', 3'', and 4''. Compound **1** showed similarities with the known compound **2** with observed variation at C-2'' (δ_{C} 86.6 for compound **1** whereas δ_{C} 76.1

for compound **2**), which was similar to δ_{C} 89.5, observed in lachnoisoflavones A,¹⁸ which suggested the cyclization in compound **1**. It was also confirmed by comparing the oxo carbon (C-4) value of compound **1** (δ_{C} 181.2) with lachnoisoflavone A (182.9 ppm). Correlated spectroscopy spectrum showed the following correlations (Figure 1): δ_{H} 7.39 (H-2', 6') with δ_{H} 6.90 (H-3', 5'), δ_{H} 6.71 (H-1''') with δ_{H} 5.61 (H-2'''), 4.45 (H-2'') with δ_{H} 3.16 (H-1''a) and δ_{H} 3.03 (H-1''b). Based on spectral data, compound **1** is identified to be a rare new natural pentacyclic isoflavonoid including a rare 5-membered dihydrofuran ring between C-5 and C-6. Finally, compound **1** was elucidated as 10-(4-hydroxyphenyl)-5,5-dimethyl-2-(prop-1-en-2-yl)-2,3-dihydro-5H,11H-furo[2,3 f]pyrano[2,3 h]chromen-11-one and named as leptoisoflavone A.

The known isolated compounds (Figure 2) were identified as euchrenone b₉ (**2**),¹⁹ auricularin (**3**),²⁰ kaempferol (**4**), avicularoside (**5**),²¹ myricetin-3-arabinoiside (**6**),²²




Figure 1. Key HMBC and COSY correlations of compound **1**.

trans-*N*-feruloyltyramine (**7**), *trans*-*N*-caffeoyltyramine (**8**),²³ and β -sitosterol (**9**).

Crude methanolic extract of aerial parts of *L. leptophyllum* exhibited moderate activity toward endocannabinoid receptors (cannabinoid receptor type 1 [CB1] and CB2). Auricularin (**3**) showed activity toward CB1 receptor (86.7% displacement with IC_{50} 8.92 μ M).¹⁵

Experimental

General

Bruker model AMX 500 NMR and 400 NMR spectrometers operating on a standard pulse system were used to acquire 1H and ^{13}C NMR and 2D spectra. The instruments ran at 500 and 400 MHz for 1H whereas they ran at 125 and 100 MHz for ^{13}C . $CDCl_3$, dimethyl sulfoxide- d_6 , and acetone- d_6 were used as NMR solvents, and trimethyl silane was used as an internal standard. Electrospray ionization-mass spectrometric (ESI-MS) data were recorded on Thermo Orbitrap Fusion (Thermo Scientific). Samples were analyzed in the positive mode of ionization and were directly infused at 3 μ L/min. Mass was analyzed in Orbitrap (mass error on the instrument <2 ppm). Electrospray ionization-MS data were obtained on a Micromass Q-ToF micromass spectrometer. Fourier transform mass spectrometry-ESI was analyzed on Thermo Orbitrap Fusion (Thermo Scientific). The sample was analyzed in the positive mode of ionization. Thin-layer chromatography was performed on precoated silica gel GF254 plates and column chromatography was performed on silica gel (200-300 mesh) and Sorbadex-LH20 (Sorbent Technologies, Atlanta, GA, USA) with detection provided by UV light (254 and 366 nm) and by spraying with 1% vanillin- H_2SO_4 reagent followed by heating for 5 to 10

minutes (105°C). Quantitative high-performance liquid chromatography (HPLC) was conducted using an Agilent 1100 HPLC system equipped with a degasser (G1379A), quaternary pump (G13311A), auto sampler (G1313A), column oven (G1316A), and UV-diode array detector (G1315B) controlled by Chemstation software. The analysis was carried out on RP-C18 columns (150 \times 4.6 mm; particle size 5 μ m; Luna) and (250 \times 10.0 mm; particle size 10 μ m; Luna) with column oven temperature set at 25°C and using the isocratic system of eluent water (A) and acetonitrile (B) for the separation of the target compounds. Acetonitrile and water solvents were of HPLC grade, where acetic acid was added as a modifier to achieve a final concentration of 0.1% in each solvent. The flow rates of the solvent were 1.0 mL/min for the analytical injections whereas 5 mL/min for the semipreparative ones and the injection volumes were 5.0 and 50 μ L for the analytical and semipreparatives, respectively. All the analysis was carried out at wavelengths of 254 and 280 nm with a run time of 40 minutes.

Plant Material

Limonium leptophyllum (Schrenk) O. Kuntze aerial parts were collected in August 2016, from Kyzylorda region, Kazakhstan. The plant material was authenticated by Akhtaeva Nursulu and the specimen voucher (No.0916.) was stored at the department of biology and biotechnology of al-Farabi Kazakh National University.

Extraction and Isolation

Limonium leptophyllum aerial parts (830 g) were extracted using MeOH (3 L \times 3 times); the extract was thereafter filtered and concentrated on a rotary evaporator at 40°C yielding 90.3 g of crude methanolic extract. This extract (50 g) was fractionated over HP 20 gel (200 g) column using H_2O -methanol (1:9), methanol, acetone, and dichloromethane (DCM) (each 750 mL). Each fraction was evaporated to dryness to give 16.47 g fraction A (H_2O :methanol 1:9), 7.87 g fraction B (methanol), 11.25 g fraction C (acetone), and 1.17 g fraction D (DCM).

Fractions C and D were combined and the mixture (E, 12 g) was loaded onto a normal phase silica gel vacuum liquid column (210 g) and eluted using hexane-acetone gradient to yield 12 fractions (E1-E12) and continued with methanol-DCM gradient to yield 10 more fractions (E13-E22). Fraction E5 was identified as β -sitosterol (**9**, 20 mg). Fraction E13 was identified as kaempferol (**4**, 10 mg). Fractions E5-E8 were combined together and purified on HPLC using water-acetonitrile gradient 60:40 to 20:80 over a period of 20 minutes, to yield euchrenone b₉ (**2**, 2.5 mg, RT 13.1 min) and compound **1** (2.1 mg, RT 14.6 min). Fractions E11 and 12 were combined (497 mg) and




Figure 2. Compounds isolated from *L. leptophyllum*.

purified over silica gel column to afford auriculasin (**3**, 25 mg). Combined fractions E20-22 (5 g) were loaded on silica gel and purified using methanol-DCM gradient system to afford avicularoside (**5**, 10 mg), myricetin-3-arabinoside (**6**, 15 mg), *trans*-*N*-feruloyltyramine (**7**, 5 mg), and *trans*-*N*-caffeoyltyramine (**8**, 8 mg).

Leptoisoflavone a (I)

Yellow solid.

$[\alpha]_D^{22.5}$: -3.125 (c 0.064, MeOH).

^1H and ^{13}C NMR: Table 1.

High-resolution ESI-MS: m/z 403.1532 $[\text{M} + \text{H}]^+$ (calc. $\text{C}_{25}\text{H}_{22}\text{O}_5 + \text{H}$, 403.1545).

Cannabinoid and Opioid Receptor Assays

The affinity of the total extracts, fractions, and isolated compounds toward cannabinoid and opioid receptors was analyzed according to the published method.²⁴ The crude methanolic extract exhibited moderate activity toward endocannabinoid receptors. Compound **3** showed activity toward CB1 receptor.

Acknowledgments

The authors thank Dr Jon Parcher for proofreading the manuscript. The authors acknowledge Award Number P20GM104932 from the National Institute of General Medical Sciences for bioassay results.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The

project was supported by National Center for Natural Product Research, School of Pharmacy, University of Mississippi, University, MS 38677, USA. This work was supported in part by the USDA Agricultural Research Service Specific Cooperative Agreement No. 58-6060-6-015.

Supplemental Material

Supplemental Material: Supplemental material for this article is available online.

References

- Zhusupova GE. *Chemical Composition of Limonium Mill Genus Plants and Creation of Preparations on its Basis [dissertation]*. Almaty, Kazakhstan: Al-Farabi Kazakh National University; 2007.
- Aniya Y, Miyagi C, Nakandakari A, Kamiya S, Imaizumi N, Ichiba T. Free radical scavenging action of the medicinal herb *Limonium wrightii* from the Okinawa islands. *Phytomedicine*. 2002;9(3):239–244.
- Kandil FE, Ahmed KM, Hussieny HA, Soliman AM. A new flavonoid from *Limonium axillare*. *Arch der Pharm*. 2000;333(8):275–277.
- Murray AP, Rodriguez S, Frontera MA, Tomas MA, Mulet MC. Antioxidant metabolites from *Limonium brasiliense* (Boiss.) Kuntze. *Zeitschrift für Naturforschung*. 2004;59(7-8):477–480.
- Kuo YC, Lin LC, Tsai WJ, Chou CJ, Kung SH, Ho YH. Samarangenin B from *Limonium sinense* suppresses herpes simplex virus type 1 replication in Vero cells by regulation of viral macromolecular synthesis. *Antimicrob Agents Chemother*. 2002;46(9):2854–2864.
- Gadetskaya AV, Mohamed SM, Tarawneh AH, et al. Phytochemical characterization and biological activity of secondary metabolites from three *Limonium* species. *Med Chem Res*. 2017;26(11):2743–2750.
- Gadetskaya AV, Tarawneh AH, Zhusupova GE, et al. Sulfated phenolic compounds from *Limonium caspium*: isolation,

- structural elucidation, and biological evaluation. *Fitoterapia*. 2015;104:80–85.
8. Medini F, Fellah H, Ksouri R, Abdely C. Total phenolic, flavonoid and tannin contents and antioxidant and antimicrobial activities of organic extracts of shoots of the plant *Limonium delicatulum*. *J Taibah Univ Sci*. 2014;8(3):216–224.
 9. Liebezeit G, Künnemann TD, Gad G. Biotechnological potential of North Sea salt marsh plants—a review of traditional knowledge. *J Biotechnol*. 1999;70(1-3):77–84.
 10. Lin LC, Chou CJ. Flavonoids and phenolics from *Limonium sinense*. *Planta Med*. 2000;66(4):382–383.
 11. Rastogi S, Pandey MM, Rawat AKS. Traditional herbs: a remedy for cardiovascular disorders. *Phytomedicine*. 2016;23(11):1082–1089.
 12. Tang XH, Yu F, Liu J, Gao J, Yan LF, Dong MM. Isolation and identification of anti-tumor polysaccharide LSP21 from *Limonium sinense* (Girard) Kuntze. *Int J Bio Macromol*. 2014;70:138–142.
 13. Yang LL, Yen KY, Kiso Y, Hikino H. Antihepatotoxic actions of Formosan plant drugs. *J Ethnopharmacol*. 1987;19(1):103–110.
 14. Zhusupova GE, Artamonova NA, Abilov ZA, Orazbaeva ZK. Lipophilic pigments and fatty acids from the aerial parts of certain plant species of the genus *Limonium*. VII. *Chem Nat Compd*. 2006;42(5):512–514.
 15. Orazbekov Y, Ibrahim MA, Mombekov S, et al. Isolation and biological evaluation of prenylated flavonoids from *Maclura pomifera*. *Evid Base Compl Alternative Med*. 2018; doi: 10.1155/2018/1370368.
 16. Satmbekova D, Srivedavyasari R, Orazbekov Y, Omarova R, Datkhayev U, Ross SA. Chemical and biological studies on *Cichorium intybus* L. *Nat Prod Res*. 2018;32(11):1343–1347.
 17. Srivedavyasari R, White MB, Kustova TS, Gemejiyeva NG, Cantrell CL, Ross SA. New tetranorlabdanoic acid from aerial parts of *Salvia aethiopis*. *Nat Prod Res*. 2018;32(1):14–17.
 18. Awouafack MD, Spiteller P, Lamshöft M, et al. Antimicrobial isopropenyl-dihydrofuranisoflavones from *Crotalaria lachnophora*. *J Nat Prod*. 2011;74(2):272–278.
 19. Mizuno M, Matsuura N, Iinuma M, Tanaka T, Phengklai C. Isoflavones from stems of *Euchresta horsfieldii*. *Phytochemistry*. 1990;29(8):2675–2677.
 20. Monache GD, Scurria R, Vitali A, et al. Two isoflavones and a flavone from the fruits of *Maclura pomifera*. *Phytochemistry*. 1994;37(3):893–898.
 21. Kalegari M, Miguel MD, Dias JdeFG, et al. Phytochemical constituents and preliminary toxicity evaluation of leaves from *Rourea induta* Planch. (Connaraceae). *Braz J Pharm Sci*. 2011;47(3):635–642.
 22. Arot LOM, Midiwo JO, Kraus W. A flavonol glycoside from *Myrsine africana* leaves. *Phytochemistry*. 1996;43(5):1107–1109.
 23. Al-Taweel AM, Perveen S, El-Shafae AM, et al. Bioactive phenolic amides from *Celtis africana*. *Molecules*. 2012;17(3):2675–2682.
 24. Tarawneh A, León F, Pettaway S, et al. Flavonoids from *Perovskia atriplicifolia* and their in vitro displacement of the respective radioligands for human opioid and cannabinoid receptors. *J Nat Prod*. 2015;78(6):1461–1465.