

University of Mississippi

eGrove

---

Electronic Theses and Dissertations

Graduate School

---

1-1-2023

## Analysis of Dietary Micronutrient Intake of NCAA Track and Field Athletes

Haley Noelle Melton

Follow this and additional works at: <https://egrove.olemiss.edu/etd>

---

### Recommended Citation

Melton, Haley Noelle, "Analysis of Dietary Micronutrient Intake of NCAA Track and Field Athletes" (2023). *Electronic Theses and Dissertations*. 2549.  
<https://egrove.olemiss.edu/etd/2549>

This Thesis is brought to you for free and open access by the Graduate School at eGrove. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of eGrove. For more information, please contact [egrove@olemiss.edu](mailto:egrove@olemiss.edu).

**ANALYSIS OF DIETARY MICRONUTRIENT INTAKE OF NCAA TRACK AND  
FIELD ATHLETES**

A Thesis  
presented in partial fulfillment of requirements  
for the degree Master of Science  
in the Department of Nutrition and Hospitality Management  
The University of Mississippi

by

HALEY N. MELTON

May 2023

Copyright Haley N. Melton 2023

ALL RIGHTS RESERVED

## ABSTRACT

Vitamins and minerals play an important role in many functions within the body and contribute to maintenance of health and well-being of all individuals, including athletes. An athlete's performance could be negatively impacted if micronutrient recommendations are not met consistently. The purpose of this study was to analyze the dietary micronutrient consumption of NCAA Division I Collegiate Track and Field athletes. Thirty-seven track and field athletes, 15 females and 22 males, participated in the study. Multiple 24-hr recalls were collected, including weekdays and weekends, and analyzed using NSDR to obtain average dietary micronutrient intakes, which could then be compared to the Recommended Daily Allowance (RDA) for each micronutrient. Pearson Chi Squared tests were used to assess correlations amongst the various participant characteristics, such as gender, age, race, track event group, and place of residence. Results indicated a significant relationship between gender and dietary iron intake ( $p < 0.001$ ), place of residence and calcium consumption ( $p < .05$ ), and the participants' race and magnesium intake ( $p < .05$ ). A cause for the high prevalence of inadequate dietary micronutrient consumption could be related to the commonness of energy deficits found in student-athletes. The prevalence of under-consuming micronutrients indicates a need for sports dietitians to focus on the assessment of vitamin and mineral dietary intakes and provide education surrounding the importance of adequate micronutrient intake to prevent negative health and performance implications associated with inadequate levels.

## LIST OF ABBREVIATIONS AND SYMBOLS

RDA Recommended Daily Allowance

EAR Estimated Average Requirement

MCG Microgram

MG Milligram

## TABLE OF CONTENTS

ABSTRACT.....	ii
LIST OF ABBREVIATIONS.....	iii
INTRODUCTION.....	1
REVIEW OF LITERATURE.....	4
METHODS.....	16
RESULTS.....	18
DISCUSSION.....	22
LIST OF REFERENCES.....	29

LIST OF TABLES

**Table 1.** List of Essential Vitamins and Minerals.....1

**Table 2.** RDA for Nutrients.....8

**Table 3.** Demographics of Participating Student-Athletes at Division I University in  
Southeastern U.S.....18

**Table 4.** Prevalence of Under Consuming Relative to the RDAs for Multiple Micronutrients  
(Excluding Vitamin D).....19

**Table 5.** Number of Deficient Participating Athletes for each Selected Micronutrient.....20

**Table 6.** Student-Athletes Under Consuming RDA for Each Micronutrient.....20

## I. INTRODUCTION

Micronutrients play a vital role in a human's overall health and well-being. Both vitamins and minerals assist in daily physiological functions such as digestion, metabolism, cellular growth, and reproduction.<sup>1,2</sup> Micronutrients may only be needed in small amounts compared to macronutrients, but serious health and performance implications can occur when the body becomes deficient or insufficient.<sup>3</sup> Many vitamins and minerals are considered essential because they cannot be obtained from the diet; therefore, it is crucial for individuals to obtain a well-balanced diet to ensure they gain the nutrients required for a healthy life.<sup>1</sup> A list of these essential micronutrients can be found in Table 1.

**TABLE 1. LIST OF ESSENTIAL VITAMINS AND MINERALS**

<u>Vitamins</u>		<u>Minerals</u>	
Vitamin A	Thiamin	Calcium	Chloride
Riboflavin	Niacin	Chromium	Copper
Vitamin B <sub>6</sub>	Folate	Fluoride	Iodine
Pantothenic Acid	Biotin	Iron	Magnesium
Choline	Vitamin B <sub>12</sub>	Manganese	Molybdenum
Vitamin C	Vitamin D	Nickel	Phosphorus
Vitamin E	Vitamin K	Potassium	Selenium
		Sodium	Zinc

One of the best ways to obtain the recommended amounts of micronutrients is through a well-balanced, nutritious diet. A variety of foods and beverages in the diet provide the body with



various different nutrients, but there is not one singular item that can provide the body with everything it requires to function properly. Instead, we must incorporate a variety of foods in order to ensure the body is able to get all of the macronutrients, vitamins, and minerals it requires to carry out its daily physiological functions.<sup>1,2,4,5</sup> Eliminating, or even restricting any of the food groups from usual intake increases an individual's risk of developing detrimental health effects resulting from nutrient deficiencies.<sup>1,4-8</sup> Along with health effects, athletes could see negative impacts on their performance when trying to compete with deficiencies of specific nutrients.

Consuming a balanced diet with sufficient variety to incorporate all nutrients is easier said than done for a lot of individuals. Difficulties arise when an individual lacks the knowledge of how to incorporate different food items to ensure they are obtaining all of the essential nutrients. Aside from adults who are pursuing a career in a nutrition-related field, many people in the United States have limited nutrition knowledge, which can negatively impact their dietary selections.<sup>3,9-11</sup> Although nutrition knowledge does not always translate to behavior, it is difficult to expect adequate dietary intakes from individuals who do not have the knowledge related to the various food groups, much less the importance of each vitamin and mineral.

The inability to obtain enough food and a well-balanced variety of items to meet estimated needs is another factor that negatively impacts an individual's ability to consume recommended amounts of nutrients. Food insecurity, insufficient amounts or quality of food to support an individual's needs, is a prevalent issue on many college campuses.<sup>12,13</sup> A study concluded 35 to 42% of college students struggle with food insecurity.<sup>12</sup> Being a student-athlete does not guarantee the individual is food secure. However, Reader et al. concluded in their study analyzing food insecurity in Division I athletes 60% of student-athletes reported food insecurity, and 37% fell into the "very low" food insecurity category.<sup>13</sup> Division I is the highest level of

collegiate athletics, yet despite the affluence of athletics programs at these institutions, athletes still have a difficult time obtaining adequate amounts of food.<sup>13,14</sup> The struggle of food insecurity could be a reason student-athletes are under consuming compared to the RDA because they are challenged to consume enough calories overall.

Current research indicates under consuming the recommended intakes for various micronutrients is prevalent in the U.S., approximately 31% of the population.<sup>2,5,6,15</sup> Track and field athletes in particular present risks of under consumption because of their lack of nutrition knowledge and increased risk of food insecurity due to the limited amount of student-athletes that receive scholarship money for this sport.<sup>16</sup> Collegiate track and field programs have upwards of 100 student-athletes on their rosters but are only provided a minimal amount of scholarships to divide among those athletes. Track and field teams assign scholarships with an equivalency system and it is rare for an individual to receive anything other than a partial scholarship, requiring the student-athlete to contribute funds to the cost of attending a college or university.<sup>16</sup>

A gap in the literature fails to include the prevalence of NCAA track and field athletes at risk of under consuming vital nutrients. The purpose of this study was to assess NCAA track and field student athlete's consumption of micronutrients. Vitamin C, vitamin D, iron, calcium, and magnesium were selected for this study primarily because of the negative performance effects associated with under consumption in athletes and prevalence of deficiency depicted in current literature. Understanding the prevalence of under consumption of essential micronutrients can create awareness and the need for education surrounding the intake of these nutrients, not only to benefit an athlete's performance, but also improve their health status in the short and long term.

## II. REVIEW OF LITERATURE

### *Prevalence and Symptomology of Deficiencies*

In order to prevent deficiencies from occurring it is recommended that individuals aim to consume recommended amounts of each nutrient daily. These values have been determined by the Food and Nutrition Board of the Institute of Medicine and are known as the Recommended Daily Allowance (RDA) and Estimated Average Requirement (EAR).<sup>17</sup> Symptoms may occur in individuals who are under consuming the recommended intakes for micronutrients, even if they are not clinically diagnosed as deficient.<sup>3</sup> These symptoms may include irritability, fatigue, soreness, impaired immune function and irregular heart rhythms.<sup>2</sup> Current research suggests the best way to prevent micronutrient deficiencies, even on a sub-clinical level, is through consistent intake of a well-balanced diet that is suitable for the individual's caloric and macronutrient needs.<sup>7</sup> Unfortunately, studies have found that a low proportion of people in the United States consume a diet that provides recommended amounts of all essential vitamins and minerals.<sup>2,5</sup> Adequate levels of micronutrients are crucial for proper health, but up to one-third of the population are at risk of micronutrient deficiencies, sometimes as a result of under consumption through the diet.<sup>2</sup> One study concluded that a majority of the U.S. population under-consumes many essential micronutrients, including vitamin D, vitamin C, calcium, and iron when compared to the EAR.<sup>2</sup> According to McClung et al., women are at a higher risk of micronutrient deficiencies than men, who are often consuming macronutrients and micronutrients at sub-optimal amounts.<sup>6,7</sup> The most common nutrient deficiencies in females was vitamin D, calcium, and iron.<sup>15</sup> All females are predisposed to a higher risk of deficiencies, stemming from a

generalization of inadequate energy availability or under consuming particular macronutrients.<sup>2,4</sup> Specifically, it seems women are more likely to have iron and vitamin D deficiencies compared to their male counterparts.<sup>2</sup> Men are not exempt from micronutrient deficiencies however. One study found a higher prevalence of vitamin C deficiency in males (8.7%) compared to females.<sup>2</sup>

Athletes are also susceptible to vitamin and mineral deficiencies. In fact, Heffernan et al. concluded that up to 60% of athletes may be deficient in some micronutrients.<sup>5</sup> The literature currently does not provide adequate evidence to conclude that sufficient levels of micronutrients, or elevated intakes of specific micronutrients, can positively impact athletic performance.<sup>5</sup> However, it can be concluded that deficient or insufficient levels of vitamins and minerals can cause detrimental effects to performance because of the symptoms associated with the deficiencies.<sup>2,3</sup>

The literature confirms the best way to avoid micronutrient deficiencies is to consume an adequate amount of calories to match an individual's energy expenditure.<sup>1,4,6</sup> This can be achieved when consuming from a wide variety of foods that supports their health status and promotes optimal performance in athletes.<sup>7</sup> Unfortunately, the primary cause of micronutrient deficiencies is related to under consumption of total energy, either unintentionally associated with a lack of knowledge, or in a restrictive manner.<sup>4,6-8,15</sup> People who are not consistently obtaining adequate calories and macronutrients have the potential to develop micronutrient deficiencies as well.<sup>7</sup> An athlete may be at a further disadvantage because of their higher energy demands associated with the high level of physical activity they frequently participate in, but it is currently unknown if athlete's have a higher demand for micronutrients and therefore increased recommended amounts.<sup>1,3-6</sup>

Under consumption of vitamins and minerals for an extended period of time can result in insufficient or deficient levels of micronutrients, causing serious health and performance implications for the individual, such as fatigue, lack of concentration, or delayed muscle recovery.<sup>2,5</sup> In the early stages of low levels some symptoms may be difficult to associate with micronutrient deficiencies because they can be linked with many different etiologies. Prolonged deficiencies can exacerbate to more serious conditions, such as decreased bone mineral density, anemia, diabetes, cardiovascular and kidney disease.<sup>2,5</sup> Micronutrient deficiencies can develop as a result of many different factors, including but not limited to under consumption of the nutrient, low energy availability, elevated demands, and difficulties with absorption or metabolism of the nutrient.<sup>1,4-6,15</sup> The compounding effects of these situations overtime can cause health and performance implications to the athlete that can reduce their time of play or potentially cease all activity completely until the problem is resolved. It is important for athletes to be aware of the potential causes and risks associated with micronutrient deficiencies in order to prevent these occurrences and increase the maximal amount of time spent playing their sport.

### ***Athletes and Micronutrient Deficiencies***

As an athlete it is common to put your body through immense periods of stress during seasons of training and competition to consistently perform at their highest potential. Athletes require high energy intakes in both calories and macronutrients to match their intense physical demand. Appropriate and adequate intake allows the athletes to continue performing at a high level while adequately recovering, reducing the risks of injuries, and progressing with their training.<sup>6-8,15,18</sup> When an athlete is consistently in an energy deficit, there is a heightened risk of detrimental health and performance consequences.<sup>3,7,15,18</sup> The literature does not confirm increased recommended values of micronutrients for athletes as it does for caloric and

macronutrient intakes, however, under consumption of essential micronutrients, particularly vitamin D, calcium, magnesium, iron and vitamin C, have also shown to have negative health and performance impacts.<sup>1</sup>

Current research indicates the selected micronutrients being considered in this study have the largest potential impact on student-athletes and emphasis is placed on adequate consumption of these nutrients within the diet.<sup>1,3-7,15</sup> Inadequate consumption of vitamin D and calcium has been associated with lowered bone mineral density and an increased risk of bone related injuries, such as stress fractures, a common injury among athletes.<sup>1,6,15,20,21</sup> Insufficient iron levels and anemia can negatively impact an athlete's performance because of inadequate oxygen being transported to the muscles during times of exercise.<sup>1,5,6</sup> Athletes may experience increased levels of fatigue and the inability to adapt to training, negatively impacting performance. Vitamin C, an enhancer for iron absorption, is another important nutrient for athletes and should be consumed in sufficient amounts.<sup>2,5,22</sup> Not only does it increase the absorption ability of dietary iron, but it also acts as an anti-inflammatory agent, improving an athlete's ability to recover by reducing prolonged inflammation within the body.<sup>1,19,22</sup> Low levels of magnesium can also inhibit recovery in athletes and should be recognized among the others as an important micronutrient in the dietary intakes of student-athletes.<sup>5,23</sup> Research is continuing to accumulate regarding magnesium's role in the body's ability to recover following training, but current research indicates adequate levels of magnesium can reduce exercise-reduced inflammation and maintain the muscles integrity, allowing an athlete quicker recovery time following intense training loads.<sup>5</sup>

#### ***Overview of Nutrients Important for Athletes:***

The following vitamins and minerals were selected because of their significance in health and performance and the significant amount of research associated with athletes and the

detrimental effects deficiencies can have on performance.<sup>1,3-7,15</sup> It is acknowledged that these are not the only nutrients that can negatively impact health and performance when under consumed, but these have significant research associated with the negative impacts at present. RDA values for each selected micronutrient is indicated in Table 2.

**TABLE 2. RDA FOR NUTRIENTS**

	<b>Females 19-30 y</b>	<b>Males 19-30 y</b>
Vitamin D	15 mcg	15 mcg
Calcium	1,000 mg	1,000 mg
Iron	18 mg	8 mg
Vitamin C	90 mg	75 mg
Magnesium	400 mg	310 mg
Abbreviations: mcg = microgram, mg=milligram		

*Vitamin D*

Vitamin D benefits the body through its role in supporting and regulating bone health and assisting with the absorption of calcium into the cells.<sup>15,20,21</sup> In addition to assisting with the regulation of calcium levels, vitamin D is also associated with maintaining skeletal muscle health and supporting our immune function.<sup>15,21</sup> Sufficient vitamin D levels are critical for all individuals, including athletes, in reducing the risk of bone injuries, such as stress fractures.<sup>6,15,20,21</sup> Vitamin D can be obtained from the diet, although foods containing the nutrient are limited.<sup>6,15,21</sup> Aside from dietary consumption of vitamin D, another way of obtaining the nutrient from UVB rays from the sun being absorbed through the skin.<sup>20,21</sup> Current research suggests that sufficient levels of vitamin D can “be obtained through 600 IU/day of dietary vitamin D, even in the absence of sun exposure” but McClung et al. believes that is an

underestimated value according to emerging research surrounding optimal vitamin D dietary intake.<sup>6</sup> Many studies have concluded that athletes are under consuming the necessary amounts of vitamin D, leading to suboptimal blood levels. Despite the vitality of having sufficient levels of vitamin D to promote bone health in athletes, it is not unusual for athletes to have insufficient or deficient levels of vitamin D because of the difficulties associated with obtaining the nutrient from the diet as well as from the sun. Vitamin D deficiency, 25(OH)D blood levels < 20ng/mL, is a concern for many athletes playing both indoor and outdoor sports.<sup>4,21</sup> One study found that 33-42% of female athletes may have insufficient blood 25(OH) D levels.<sup>6</sup> Other studies have concluded vitamin D insufficiency ranges from 13.3%-80% and vitamin D deficiency rates of .05%-72.8% in various athletes.<sup>4,21</sup> Specific factors can contribute to an athletes risk of developing vitamin D deficiency, including time of year, location, darker skin pigmentation, amount of time spent outside, and skin exposure from clothing worn.<sup>6,21</sup> With sun absorption being a primary contributor to vitamin D status, dark pigmented skin tones are reported to have a higher risk of developing vitamin D deficiencies because the melatonin levels in the skin interfere with the absorption rates of the UVB rays.<sup>4,21</sup> Athletes who identify with these factors should be cautious of the increased risk they present for low vitamin D status.

### *Calcium*

Along with vitamin D, calcium plays a vital role in the development and maintenance of bone health.<sup>6</sup> Without sufficient levels of calcium the bone integrity can disintegrate, leading to serious implications like stress fractures or decline in bone density.<sup>1,6,24</sup> Reduced bone mineral density can develop into serious medical conditions like rickets in children and osteomalacia in adults.<sup>6</sup> The micronutrients impact on bone health is so essential that in 2010 the Institute of Medicine (IOM) took the initiative to update the dietary reference intake levels of dietary



calcium and vitamin D with bone health as the main motivating factor in their decision. Other health contributions associated with the nutrient did not have sufficient amounts of research to support the credibility like bone health.<sup>6</sup> Unfortunately, similar to vitamin D, calcium intake in the general population and among athletes does not meet the recommended values which is 1,000 mg for both males and females ages 19-50.<sup>4</sup> According the NHANES data, both males and females are guilty of under consuming the recommended amounts of dietary calcium.<sup>6</sup> Contrary to vitamin D, calcium can be found in more foods, most commonly in dairy products. Under consumption of calcium-rich foods can lead to long term health implications, like osteomalacia or osteoporosis.

### *Iron*

Iron, an essential mineral that must be obtained from the diet, contributes a great deal to athletic performance and overall health. The primary role for iron includes supporting the production of proteins, known as hemoglobin and myoglobin, in the transportation and storage of oxygen to different regions within the body.<sup>6</sup> The easiest way to absorb iron is through the consumption of heme iron, which is abundant in animal products including red meat, poultry, and seafood. The other type of dietary iron is nonheme iron and is found in plant products, and is not as readily available or absorbable as heme iron.<sup>6</sup> Other nutrients associated with non-heme iron foods, like phytates and phenolic compounds interfere with absorption.<sup>6</sup> Adequate amounts of dietary iron to support sufficient blood levels can be obtained from a diet containing “highly bioavailable” sources of heme and non-heme iron in accordance with nutrients known to enhance iron absorption. Foods rich with ascorbic acid, found in citrus fruits and leafy green vegetables enhance iron absorption and are essential in obtaining satisfactory iron levels.<sup>6</sup> The RDA for iron varies for females and males. It is recommended for females 18 mg/day and males 8 mg/day,

because of monthly blood losses associated with menstruation.<sup>6</sup> One article determined that 67% of athletes do not meet the RDA values of iron.<sup>4</sup> Iron deficiency is common among athletes, reported 4-50% of male athletes and 20-50% of female athletes, but some studies suggesting the prevalence is above 50%.<sup>3,15</sup>

There are multiple contributing factors to affect iron levels within the body. Low overall intake, low animal protein consumption, under consumption of iron-rich foods, and external factors contribute to low iron levels in the body.<sup>4-6</sup> Athletes have an elevated risk of developing deficiency or anemia because of the difficulty iron has being absorbed following intense exercise. Acute inflammation as a response to physical activity increases the levels of hepcidin circulating in the blood. Hepcidin, a hormone that regulates iron levels in the blood, becomes elevated and reduces iron's absorption rate, contributing to decreased iron stores.<sup>6,7</sup> Iron losses associated with hemolysis from repetitive ground strikes in endurance athletes and/or gastrointestinal bleeding following intense exercise is another hinderance for athletes to combat when maintain adequate iron stores.<sup>6,15</sup> Female athletes have an additional barrier for iron stores because of monthly blood losses from menstruation, which provides major iron losses via excretion in the blood, especially for those athletes with heavy menstrual cycles.<sup>6,7,15</sup> Any athlete consuming a restrictive diet, especially limiting sources of heme iron, increases the risk of iron deficiency.<sup>7,15</sup> Unfortunately for athletes, the most prevalent symptom associated with insufficient blood levels of iron is decline in athletic performance due to the lack of oxygen carrying capacity from the red blood cells.<sup>6,15</sup> This can be detrimental to athletic performance, reducing endurance capacity and inducing muscle fatigue.<sup>5,6</sup> Iron acts as a cofactor in the brain as well as the skeletal muscles, so decline in concentration, behavior, and cognitive function is also associated with iron deficiency.<sup>6</sup>

### *Vitamin C*

Vitamin C is another micronutrient that cannot be synthesized within the body and must therefore be consumed as part of the diet.<sup>19</sup> In the body vitamin C contributes to many functions, including collagen tissue synthesis, absorption of iron from the blood stream, synthesis of hormones, and acts as an antioxidant to reduce reactive oxygen species (ROS) that could potentially cause harm to tissue.<sup>19,22</sup> For athletes it is important to consume recommended amounts of vitamin C daily to maintain the redox balance and reduce inflammation following physical activity. The RDA for vitamin C is 75 mg/day for women and 90 mg/day for men.<sup>17</sup> Regular consumption of fruits and vegetables accounts for a majority of a person's daily intake of vitamin C.<sup>19</sup> According to Jordan et al. the nutrient is found in a variety of fruits and vegetables and it is uncommon for athletes to consume less than the RDA.<sup>4</sup> Controversially, Bird et al. found deficiency of vitamin C in 8.7% of adult males aged 19-50 years and contributed this finding to the lack of fruits and vegetables selected for consumption in their diet.<sup>2</sup> The article also associated the varying motivational factors for incorporating fruit and vegetables into the diet as a "gender-based difference".<sup>2</sup> Under consumption of vitamin C can lead to poor recovery, inhibiting future performance in athletes and should aim to be regularly consumed in the diet.

### *Magnesium*

Many functions that occur within the human body are reliant on magnesium to regulate processes and function as a cofactor for the numerous enzymes involved. A function of significance includes acting as an electrolyte or an electrical charge associated with skeletal and myocardial tissue contractions. Electrolytes act to regulate muscle function during exercise, prevent muscle cramps, and can regulate blood pressure.<sup>23</sup> Magnesium also functions to support energy metabolism and emerging evidence shows that it may contribute to the maintenance of

muscle mass and systemic inflammation, all important factors for athletes to be aware of.<sup>5,23</sup> The RDA for magnesium is 400 mg for males and 310 mg for females, with approximately 10% of our magnesium requirement coming from our drinking water.<sup>17,23</sup> Despite the contribution stemming from water, many people are not obtaining the recommended daily intake values of magnesium likely leading to insufficient or deficient levels.<sup>4,23</sup> When deficiencies are present a widespread amount of diseases can develop, including hypertension, coronary artery disease, stroke, heart failure, chronic obstructive pulmonary disease, among others.<sup>23</sup> Limited research is available assessing the intake of athletes, but deficiencies could negatively impact their ability to recover and adaptation to training.

### ***Dietary Analysis***

Accurately recording participants dietary intake is a difficult task because of the many variables associated with the process and should be addressed when collecting data.<sup>25</sup> Dietary recalls rely on the participant to recollect and accurately depict their previous meals, leading to unintended reporting errors. These inconsistencies must be considered when deciding the manner in which to collect recalls.<sup>25-27</sup> Of the multiple methods for recording dietary intake, two main categories exist: real-time recording or methods of recall.<sup>25</sup> Real-time recording allows the participant to keep a food diary and record each food or beverage consumed in the moment it is ingested.<sup>25</sup> The alternative to real-time recording is methods of recall, which relies on the participants memory to accurately depict their dietary intake.<sup>25</sup> Food frequency questionnaires are one method for collecting dietary intake data. This method requires the participant to determine how frequently they consume each of the listed foods groups.<sup>25</sup> The food groups can be broad categories or list specific food items, depending on the assessment. Difficulties arise

when the food frequency questionnaire is relatively short in length and underestimates the variety of a participant's diet, unable to accurately depict a usual intake. However, if a long food frequency questionnaire is provided to participants, they could experience fatigue and rush through the questionnaire, again, reducing the accuracy of the diet reflected from the questionnaire.<sup>25</sup>

An alternative form is the 24-hour dietary recall, which assesses a participant's daily intake from a 24-hour time frame, typically the previous day. The intention for a 24-hour recall is to gather information regarding the participant's typical intake on a daily basis.<sup>25,26</sup> It is likely that one day does not accurately depict an individual's usually daily intake, so it is recommended multiple recalls be collected to obtain a better representation of the usual intake. This method of dietary analysis is the standard for data collection, and current research suggests using at least three recalls for each participant, with some researchers suggesting using upwards of 8-32 recalls.<sup>25,26</sup> One drawback of using this many recalls is reliance on the participant and ensuring they maintain interest in providing detailed descriptions of their daily intake. It should be a goal of the research to prioritize their participant's interest when determining the number of recalls to perform in hopes of limiting participant fatigue.<sup>26</sup> Another limitation associated with 24-hour recalls is the difficulty for untrained individuals to accurately depict their true dietary intake when estimating the amounts of each food category they consume.<sup>25,27</sup> The participants may or may not understand what a serving size is for each item, reducing the accuracy and reliability when comparing the recalls against each other or to recommended standard. Both food frequency questionnaires and 24-hour recalls pose the risk of recall bias.<sup>25</sup> This occurs when participants intentionally refrain from reporting their complete diet history or adjust aspects of their recall, causing it to become inaccurate. It can be difficult to account for these types of errors, but it is

encouraged by all researchers to limit these occurrences as possible. Using a consistent, uniform approach for obtaining each participant's recall can limit the amount of variability between each assessment, strengthening the reliability of the collected data.

The purpose of the study was to analyze the dietary intake of track and field student athletes and assess if their usual consumption meets the RDA standards for each of the selected micronutrients, reducing their risks of adverse health effects and declines in performance.

### III. METHODS

#### *Participants*

All participating track and field athletes (n=37) were recruited from a NCAA DI university within the Southeastern Conference. The study included athletes from the track and field event groups jumps, sprints, pole vault, and multiple event athletes and excluded distance or throw athletes. Approval was obtained from the University of Mississippi Institutional Review Board (IRB) and procedures were conducted in accordance with the principles associated.

#### *Procedures*

##### *Dietary assessment*

For the dietary analysis of each participant, 24-hour dietary recalls were used to obtain accurate depictions of the athlete's intakes. This method was selected because of the ease of obtaining an accurate depiction of a daily routine without putting a burden on the participants with keeping up accurate recording of multiple day food logs. Multiple dietary recalls were conducted for each student-athlete participating in the study, including weekdays and weekends to gain an accurate representation of their daily intake during the off-season.<sup>25</sup> While collecting the information from the participants, the "hand method" was used to help the student-athletes accurately describe the portion sizes they consume at each meal.<sup>27</sup> The participants may or may not understand what a serving size is for each item, reducing the accuracy and reliability to compare the recalls against each other or a recommended standard. Using a consistent, uniform approach for obtaining each participant's recall can limit the amount of variability between each assessment. The recalls analyzed using Nutritional Data System for Research (NDSR) software

version 50, (2019). The student-athletes micronutrient consumption and total calorie intake were collected for comparison to the RDA for each nutrient.

### ***Statistical Analysis***

Statistical analyses were conducted using SPSS statistical software version 29 and Microsoft Excel. Descriptive statistics were conducted to determine participant characteristics, micronutrient intake, and total calories. Chi-square tests were conducted to determine any significant correlations between micronutrient intake and participants demographic variables. Statistical significance was set at  $p < 0.05$ .



#### IV. RESULTS

Of the 37 student-athletes (age: 20±1 years), 15 were females and 22 were males.

Participants belonged to one of four track and field event groups: jumps, sprints, pole vault, or multi (decathletes and heptathletes). Characteristics of the participants can be seen in Table 3.

**TABLE 3. DEMOGRAPHICS OF PARTICIPATING STUDENT-ATHLETES**

<b>Participant Characteristic</b>		<b>N (%)</b>
Gender	Female	15 (41%)
	Male	22 (59%)
Age	18	2 (5%)
	19	14 (38%)
	20	9 (24%)
	21	7 (19%)
	22	4 (11%)
	23	1 (3%)
Event	Jumps	10 (27%)
	Sprints	12 (32%)
	Multi	8 (22%)
	Pole Vault	7 (19%)
Race	Black	20 (54%)
	White	16 (43%)
	Other	1 (3%)
Year of Eligibility	Freshman	5 (14%)
	Sophomore	13 (35%)
	Junior	9 (24%)
	Senior	10 (27%)
Residence	Off-Campus	30 (81%)
	On-Campus	7 (19%)

Every student-athlete was under consuming at least one of the selected micronutrients, with many athletes failing to meet the RDA for multiple micronutrients. Only one student-athlete consumed the recommended amount of dietary vitamin D. Despite the lack of prevalence in foods, vitamin D has an abundance of availability coming from the sun’s rays that could provide an athlete with substantial amounts to prevent insufficient or deficient levels. Removing vitamin D intakes from the analysis, 92% (n=34) of student-athletes were still under consuming at least one of the selected micronutrients when compared to the RDA. Table 4 categorizes the participants into separate groups, representing the number of selected micronutrients each athlete under-consumed compared to the recommendations.

**TABLE 4. PREVALENCE OF UNDER CONSUMING RELATIVE TO THE RDAS FOR MULTIPLE MICRONUTRIENTS (EXCLUDING VITAMIN D)**

<u>Number of Deficiencies</u>					
	<b>0</b>	<b>1</b>	<b>2</b>	<b>3</b>	<b>4</b>
<b>Females</b>	0 (0%)	1 (7%)	3 (20%)	3 (20%)	8 (53%)
<b>Males</b>	3 (14%)	3 (14%)	5 (23%)	11 (50%)	0 (0%)

Results from the descriptive analysis portrayed some significant correlations among participant characteristics and under consumption of each micronutrient. Chart 1 and Table 5 depict the number of participants under-consuming each micronutrient separated by gender. Four out of the five micronutrients did not have significant results comparing adequate intakes to each gender. Iron, however, did show a significant relationship with 100% of the male participants consuming the recommended dietary iron amounts, while only one female participant met the recommendation,  $X^2(1, N = 37) = 33.03, p < 0.001$ .

**Table 5. NUMBER OF DEFICIENT PARTICIPATING ATHLETES FOR EACH SELECTED MICRONUTRIENT**

	Females	Males	Total
Calcium	13 (87%)	14 (64%)	27 (73%)
Iron	14 (93%)	0 (0%)	14 (37%)
Magnesium	12 (80%)	18 (82%)	30 (81%)
Vitamin C	9 (60%)	14 (64%)	23 (62%)
Vitamin D	15 (100%)	21 (95%)	36 (97%)

The most common micronutrient to be under consumed by all participants was magnesium at 81% (n=30), followed closely by calcium at 73% (n=27). There was a significant difference in the magnesium consumption amongst Black and White participants,  $X^2 (2, N = 37) = 6.360, p = 0.042$ , with more White participants consuming the recommended amounts of magnesium compared to Black participants. One out of 20 Black participants met the recommendations (5%) while 38% of White participants consumed adequate amounts of magnesium. Additionally, there was a significant association between the consumption of calcium and living on-campus,  $X^2 (1, N = 37) = 3.970, p = 0.046$  (Table 6).

**Table 6. STUDENT-ATHLETES UNDER CONSUMING RDA FOR EACH MICRONUTRIENT**

<b>Vitamin D</b>	<b>Pearson Chi-Squared</b>	<b>P value</b>
Gender	1.507	.220
Age	4.405	.493
Event	2.141	.544
Race	.874	.646
Year of Eligibility	2.775	.428
Residence	.240	.624
<b>Calcium</b>	<b>Pearson Chi-Squared</b>	<b>P value</b>

Gender	2.399	.121
Age	12.372	.030
Event	3.433	.330
Race	4.106	.128
Year of Eligibility	8.792	.032
Residence	3.970	.046
<b>Iron</b>		
	<b>Pearson Chi-Squared</b>	<b>P value</b>
Gender	33.032	<.001
Age	5.569	.350
Event	.943	.815
Race	1.712	.425
Year of Eligibility	2.174	.537
Residence	2.036	.154
<b>Vitamin C</b>		
	<b>Pearson Chi-Squared</b>	<b>P value</b>
Gender	.05	.823
Age	2.684	.749
Event	2.537	.469
Race	4.316	.116
Year of Eligibility	2.435	.487
Residence	1.368	.242
<b>Magnesium</b>		
	<b>Pearson Chi-Squared</b>	<b>P value</b>
Gender	.019	.890
Age	15.839	.007
Event	4.203	.240
Race	6.360	.042
Year of Eligibility	2.137	.544
Residence	.524	.469

Nearly every athlete living on-campus met the recommended intake for calcium, while upper-classmen, who mainly live off-campus, fell into the inadequate range. Aside from the previously mentioned, no other characteristics showed significant differences in micronutrient intakes and consumption of the selected micronutrients was evenly dispersed between each event group.

## V. DISCUSSION

The purpose of this study was to determine the prevalence of student-athletes under consuming micronutrients compared to the RDA values, despite these nutrients being vital for their performance. Athletes failing to meet dietary intake recommendations for micronutrients does not necessarily mean they will certainly develop biological deficiencies, but it is a noteworthy associated risk.<sup>5</sup> The data collected shows 100% of participants were under-consuming at least one of the selected micronutrients. Some nutrients were consumed at a higher rate, with some participants meeting the RDA, while others were significantly under consumed. For example, 100% of males consumed the recommended amount of iron, while only 19% of all participants consumed the RDA for magnesium. The results conclude that participants who under-consume one micronutrient are likely to under-consume multiple of the selected micronutrients, with 92% (n=34) under consuming at least two of the selected vitamins and minerals. Fifty-three percent (n=8) of the women under-consumed all five micronutrients selected for the study. The male participants did not have any individuals failing to meet the recommendations for all five of the micronutrients, but 50% (n=11) fell below the RDA for 4 out of the 5 vitamins and minerals.

Similar to the literature previously mentioned, the under-consumption of micronutrients can be attributed to under-consumption of total calories.<sup>4,6,8,15</sup> Individual estimated energy recommendations for each student-athlete were unable to be obtained during this study but compared to the 2,000-calorie recommendation for the general population, thirty-two percent of these student athletes are failing to meet their estimated energy needs. This is likely an

underestimate for calorie needs if you consider the assumption that majority of collegiate, Division I student-athletes have higher energy demands than the generalized 2,000-calorie predictor, indicating a higher percentage of participants consuming low energy intakes. Individuals under consuming their recommended caloric intake limits the availability of micronutrients within their daily intake, putting each athlete at a disadvantage for obtaining the essential micronutrients from their diet and increasing the risk of adverse effects associated with micronutrient deficiencies. Students living on campus tend to have an advantage because of the accessibility to dining halls and restaurants universities strive to acquire. Despite the abundance and convenience of food on campus, 86% (6 out of 7) of students living on campus still were under consuming at least one micronutrient, primarily magnesium, where all but one student met the RDA. The other participant characteristics were not notable when comparing intakes.

#### *Vitamin D*

Of the thirty-seven participants, only one consumed the recommended amount of dietary vitamin D. This was not shocking considering the lack of vitamin D-rich foods and the availability of vitamin D coming from UVB rays. Despite the limited amounts of vitamin D-rich foods, dietary vitamin D still has a significant place in the diet of student-athletes, as well as the general population, because of the difficulties associated with obtaining an ample amount of vitamin D from the sun to promote adequate blood levels, a vital component for many functions within the body. The study's population displayed comparable results of dietary under-consumption compared to similar studies assessing vitamin D dietary intake, approximately 2.7% and 5% of college athletes obtaining the RDA from dietary sources alone.<sup>21,24</sup> There is even debate that the current RDA for vitamin D is low compared to actual needs, especially when some populations have difficulties with absorbing adequate amounts of vitamin D from the

sun's rays. For example, individuals with darker pigmentation, indoor sport athletes, those living in northern regions of the country and the amount of clothing or SPF worn outside.<sup>6,21</sup> The impact vitamin D has on many functions within the body, primarily its role regulating bone health, warrants special attention when analyzing an individual's intake and focusing on obtaining adequate levels of vitamin D.<sup>5,20,21</sup> When dietary intake is limited and factors that inhibit absorption from UVB rays, supplementation should be considered to bridge the gap for these student-athletes at risk of deficiency.

### *Calcium*

The prevalence of under-consuming calcium was lower compared to vitamin D, but still 73% of participants fell short of consuming the RDA. Males consumed calcium at a higher rate compared to their female counterparts, with 64% under-consuming compared to 87% of females not meeting the recommendation. This finding aligned with other studies findings where 51%-92% of female athletes failed to consume the RDA for calcium.<sup>5,24,28</sup>

The results from the collected data indicated that student-athletes living on-campus consumed more dietary calcium on average compared to those students living off-campus. These findings were opposite of initial assumptions that older students, the upper classmen electing to move off-campus, would consume higher amounts of calcium compared to the freshman, who were required to live on-campus. More exposure to performance nutrition and experience living on their own leads to the assumption that students living off-campus, who are able to make more of their own decisions and purchase their own groceries, would be consuming more calcium, but the results indicated otherwise. One reason we could conclude for this finding is the convenience of calcium-rich foods on campus, via dining halls, restaurants, or convenience stores that was included in the students dining plan.

Emphasizing calcium-rich foods in the diet should be a standard practice for those working with student-athletes to instill the importance of the nutrient on their health and to help them recognize how calcium can be underconsumed if not intentionally being incorporated into a daily routine. Even though calcium is more prevalent in foods compared to vitamin D, a majority of those prospective foods are dairy products.<sup>24</sup> Student-athletes who refrain from eating dairy, either medically or by preference, need to understand they could be missing calcium from their diet and learn additional ways of incorporating sufficient amounts to ensure proper bone health and reduce the risk of related injuries.<sup>24</sup>

### *Iron*

There was a significant difference of dietary iron consumption from male to female participants, with 100% of males consuming the RDA while only 7% of female participants met the recommendation. When looking at the average iron intakes, all participants consumed similar amounts with a mean intake of  $15.78 \pm 0.76$  mg/d. Females have a higher RDA for iron compared to men, 18 mg/d versus 8 mg/d, and because of that it led to an overwhelming majority of females under-consuming dietary iron and put females at a higher risk of developing insufficient iron levels or anemia.<sup>6,29</sup> An article by Weight et al. concluded similar results from their study analyzing dietary iron intakes in male and female athletes. From their cohorts, the male's average dietary iron intake met the RDA, averaging 14.8 mg/day, but the female's average failed to meet their respective RDA, averaging below 18 mg/day.<sup>30</sup> Another study assessing the dietary intakes among female marathon runners concluded that prior to their studies intervention, none of the female athletes met the prospective RDA.<sup>31</sup> The article findings, along with the present study indicates an increased emphasis of nutrition education for female athletes regarding the difficulty of reaching the recommended intake for iron. Many things inhibit the



absorption of iron into the body, including increased hepcidin levels following exercise, other mineral rich-foods, and compounds found in plant foods like phytates and phenolic compounds.<sup>6</sup> The bioavailability of iron, or lack thereof, places another obstacle for athletes and further emphasizes the importance of proper dietary intake of iron.<sup>6</sup>

### *Vitamin C*

The results from the analysis shows 62% of all athletes failed to meet the recommendation for dietary intake according to the RDA, with 60% of females and 64% of males under-consuming. These percentages are higher than a majority of the literature, which suggests inadequate consumption of vitamin C is not common among athletes or the general population.<sup>4,32</sup> However, an article from 2017 concluded 49% of participants did not meet the recommended intake value when assessing dietary consumption compared to the EAR.<sup>2</sup>

Socioeconomic status is a potential contributor to dietary vitamin C intake levels and could be a factor in the increased prevalence of under-consuming the RDA in this study's population.<sup>19</sup> A lack of nutrition knowledge encompassing the benefits of vitamin C, not only on health but in different roles of performance, could be a limitation for these student-athletes failing to consume adequate amounts of dietary vitamin C.

### *Magnesium*

Magnesium was the most under-consumed nutrient from the data with 81% of participants failing to meet the RDA for magnesium. There were no significant differences among the participant characteristic groups. One finding of interest was the group of seven participants that met the RDA for magnesium had higher total caloric intakes than the other participants, supporting the notion that consuming a diet that meets energy needs can provide student-athletes with the recommended micronutrient needs.<sup>4,5,15</sup> Other studies have concluded

magnesium to be a difficult nutrient to obtain the RDA in most athletes diet.<sup>4,5</sup> Nutrition education surrounding magnesium rich foods is essential for student-athletes to ensure they are maintaining well-rounded, substantial diets to provide them the micronutrients they need to maintain a healthy lifestyle and perform at their highest potential.

### ***Application of Results***

The results from this study indicate the need for diet education for student-athletes and the ability for them to have their diets analyzed for the under consumption of essential micronutrients. The impact these nutrients have on their performance is enough to warrant these actions, but the detrimental impacts on overall health further indicate the need for trained professionals working in athletics to further analyze student-athletes' dietary intakes because of the prevalence on under-consumption of micronutrients. Other athletic programs with similar structures should consider analyzing the micronutrient intakes of their student-athletes in addition to total caloric and macronutrient intake. Student-athletes that have limited access to dining halls, training tables, or other resources should be a primary focus because they are a primary risk for under consumption of micronutrients. Food insecurity can be prevalent on any collegiate campus and can include student-athletes with limited scholarship funding to cover meal plans. Track and field athletes tend to fall into these categories, requiring diligence when assessing the quality of their micronutrient and overall intake.

### ***Strengths and Limitations***

Using multiple 24-hour dietary recalls provided a solid representation of each participant usual dietary intake and allowed for accurate estimations of their micronutrient consumption.

Alternative methods of dietary analysis could risk increased room for error and uncertainty when analyzing the caloric and micronutrient intakes of the participants. Each recall was conducted or analyzed by a Registered Dietitian. The software used for dietary analysis had a data base of many different foods, including name branded, packaging, and common fast food menu items, which eliminated estimations and uncertainties when logging these items and provided a more accurate depiction.

While lengths were taken to minimize any limitations associated with this study, there are some that could not be avoided. Any form of dietary analysis contains some limitations because it relies on the memory of each student-athlete and their ability to accurately describe their intakes. These student-athletes do not have a background in nutrition and struggle to accurately depict a portion size for a meal. Estimations were made to best determine accurate intakes, including incorporating the “hand method” for a more accurate depiction, but room for error must be considered when analyzing their caloric and nutrient results. Blood tests could not be obtained during this study, but future research would do well to obtain blood results to confirm or reject the idea that under consumption of micronutrients from the diet leads to deficiencies with the micronutrients, increasing the potential of experiencing these negative performance consequences as well as detriments to the student-athletes health. This would have been especially beneficial for vitamin D in particular to determine if the under consumption of vitamin D from food alone accurately represents detrimental health effects even though the most abundant source of obtaining vitamin D is from UVB rays from the sun. Blood results of the nutrients would be another way to confirm the accuracy from the diet recalls in representing the athlete’s actual intakes.

## ***Conclusions***

This research study provides information regarding the frequent under-consumption of essential micronutrients from track and field student-athletes that impact both health and performance. Recognizing these tendencies can be influential when providing nutrition education to this population and encourage sports dietitian to analyze not only the caloric and macronutrient intakes of these student-athletes, but also be aware of their micronutrient intakes as well. Prioritizing the health of student-athletes is essential to promote optimal athletic performance. Analyzing micronutrient intake and screening for deficiencies can be another avenue to promote exceptional care to individuals.

## LIST OF REFERENCES

## LIST OF REFERENCES

1. Beck KL, von Hurst PR, O'Brien WJ, Badenhorst CE. Micronutrients and athletic performance: A review. *Food Chem Toxicol.* 2021;158:112618. doi:10.1016/j.fct.2021.112618
2. Bird JK, Murphy RA, Ciappio ED, McBurney MI. Risk of Deficiency in Multiple Concurrent Micronutrients in Children and Adults in the United States. *Nutrients.* 2017;9(7). doi:10.3390/nu9070655
3. Riviere AJ, Leach R, Mann H, et al. Nutrition Knowledge of Collegiate Athletes in the United States and the Impact of Sports Dietitians on Related Outcomes: A Narrative Review. *Nutrients.* 2021;13(6). doi:10.3390/nu13061772
4. Jordan S, Albracht-Schulte K, Robert-McComb J. Micronutrient deficiency in athletes and inefficiency of supplementation: Is low energy availability a culprit? 2020. 14.
5. Heffernan SM, Horner K, De Vito G, Conway GE. The Role of Mineral and Trace Element Supplementation in Exercise and Athletic Performance: A Systematic Review. *Nutrients.* 2019;11(3). doi:10.3390/nu11030696
6. McClung JP, Gaffney-Stomberg E, Lee JJ. Female athletes: A population at risk of vitamin and mineral deficiencies affecting health and performance. *ISTERH X Conf TOKYO TRACE Elem Res Health Dis.* 2014;28(4):388-392. doi:10.1016/j.jtemb.2014.06.022
7. Gastrich MD, Quick V, Bachmann G, Moriarty AM. Nutritional Risks Among Female Athletes. *J Womens Health.* 2020;29(5):693-702. doi:10.1089/jwh.2019.8180
8. Kontele I, Vassilakou T. Nutritional Risks among Adolescent Athletes with Disordered Eating. *Children.* 2021;8(8). doi:10.3390/children8080715

9. Kuczmarski M, Adams E, Cotugna N, et al. Health Literacy and Education Predict Nutrient Quality of Diet of Socioeconomically Diverse, Urban Adults. *J Epidemiol Prev Med.* 2016;2:115.
10. Yahia N, Brown CA, Rapley M, Chung M. Level of nutrition knowledge and its association with fat consumption among college students. *BMC Public Health.* 2016;16(1):1047. doi:10.1186/s12889-016-3728-z
11. Taylor M, Sullivan D, Ellerbeck E, Gajewski B, Gibbs H. Nutrition literacy predicts adherence to healthy/unhealthy diet patterns in adults with a nutrition-related chronic condition Public Health Nutrition How does Cambridge Core Share work? *Public Health Nutr.* 2019;22. doi:10.1017/S1368980019001289
12. Barry MR, Sonnevile KR, Leung CW. Students with Food Insecurity Are More Likely to Screen Positive for an Eating Disorder at a Large, Public University in the Midwest. *J Acad Nutr Diet.* 2021;121(6):1115-1124. doi:10.1016/j.jand.2021.01.025
13. Reader J, Gordon B, Christensen N. Food Insecurity among a Cohort of Division I Student-Athletes. *Nutrients.* 2022;14(21). doi:10.3390/nu14214703
14. Mayeux W, Camel S, Douglas C. Prevalence of Food Insecurity in Collegiate Athletes Warrants Unique Solutions. *Curr Dev Nutr.* 2020;4(Supplement\_2):239-239. doi:10.1093/cdn/nzaa043\_090
15. Holtzman B, Ackerman KE. Recommendations and Nutritional Considerations for Female Athletes: Health and Performance. *Sports Med.* 2021;51(1):43-57. doi:10.1007/s40279-021-01508-8

16. Barbara J. Osborne. Failing to Fund Fairly: Title IX Athletics Scholarships Compliance. *Tenn J Race Gend Soc Justice*. 2017;6(1).
17. Mahan LK, Raymond J. *Krause's Food & The Nutrition Care Process*. 14th ed. Elsevier; 2017.
18. De Souza MJ, Strock N, Ricker E, et al. The Path Towards Progress: A Critical Review to Advance the Science of the Female and Male Athlete Triad and Relative Energy Deficiency in Sport. *Sports Med*. 2022;52. doi:10.1007/s40279-021-01568-w
19. Doseděl M, Jirkovský E, Macáková K, et al. Vitamin C—Sources, Physiological Role, Kinetics, Deficiency, Use, Toxicity, and Determination. *Nutrients*. 2021;13(2). doi:10.3390/nu13020615
20. Laird E, Ward M, McSorley E, Strain JJ, Wallace J. Vitamin D and Bone Health; Potential Mechanisms. *Nutrients*. 2010;2(7):693-724. doi:10.3390/nu2070693
21. Yagüe M de la P, Yurrita LC, Cabañas MJC, Cenzual MAC. Role of Vitamin D in Athletes and Their Performance: Current Concepts and New Trends. *Nutrients*. 2020;12.
22. Kaźmierczak-Barańska J, Boguszewska K, Adamus-Grabicka A, Karwowski BT. Two Faces of Vitamin C—Antioxidative and Pro-Oxidative Agent. *Nutrients*. 2020;12(5). doi:10.3390/nu12051501
23. Alawi A, Majoni S, Falhammar H. Magnesium and Human Health: Perspectives and Research Directions. *Int J Endocrinol*. 2018;2018. doi:10.1155/2018/9041694
24. Wrzosek M, Woźniak J, Kozioł-Kaczorek D, Włodarek D. The Assessment of the Supply of Calcium and Vitamin D in the Diet of Women Regularly Practicing Sport. *J Osteoporos*. 2019;2019:1-7. doi:10.1155/2019/9214926



25. Naska A, Lagiou A, Lagiou P. Dietary assessment methods in epidemiological research: Current state of the art and future prospects. *F1000Research*. 2017;6:926.  
doi:10.12688/f1000research.10703.1
26. St. George S, Van Horn ML, Lawman H, Wilson D. Reliability of 24-Hour Dietary Recalls as a Measure of Diet in African-American Youth. *J Acad Nutr Diet*. 2016;116.  
doi:10.1016/j.jand.2016.05.011
27. Gibson A, Hsu M, Rangan A, et al. Accuracy of hands v. household measures as portion size estimation AIDS. *J Nutr Sci*. 2016;5. doi:10.1017/jns.2016.22
28. Ward KD, Hunt KM, Berg MB, et al. Reliability and validity of a brief questionnaire to assess calcium intake in female collegiate athletes. *Int J Sport Nutr Exerc Metab*. 2004;14(2):209-221.
29. Hughes RL, Holscher HD. Fueling Gut Microbes: A Review of the Interaction between Diet, Exercise, and the Gut Microbiota in Athletes. *Adv Nutr*. 2021;12(6):2190-2215.
30. Weight LM, Jacobs P, Noakes TD. Dietary iron deficiency and sports anaemia. *Br J Nutr*. 1992;68(1):253-260. doi:10.1079/BJN19920082
31. Alaunyte I, Stojceska V, Plunkett A, Derbyshire E. Dietary iron intervention using a staple food product for improvement of iron status in female runners. *J Int Soc Sports Nutr*. 2014;11:50. doi:10.1186/s12970-014-0050-y
32. de Sousa EF, Da Costa THM, Nogueira JAD, Vivaldi LJ. Assessment of nutrient and water intake among adolescents from sports federations in the Federal District, Brazil. *Br J Nutr*. 2008;99(6):1275-1283. doi:10.1017/S0007114507864841

## VITA

### EDUCATION

- Bachelor of Science (May 2020) in Nutrition Dietetics, Auburn University, Auburn, Alabama
- Dietetic Internship (May 2021) Lipscomb University, Nashville, Tennessee

### ACADEMIC EMPLOYMENT

- Graduate Assistant, Department of Nutrition and Hospitality Management, Sports Nutrition, University of Mississippi, August 2021 – August 2022. Responsibilities include providing nutrition education to Ole Miss student-athletes in the form of individual consults or group education talks. Also responsible for ordering, stocking, and maintenance of multiple fueling stations on-campus available for student-athletes.

### PROFESSIONAL MEMBERSHIPS

- Academy of Nutrition and Dietetics
- Collegiate and Professional Sports Dietitians Association